

High genetic structuring of Tula hantavirus

**Sabrina Schmidt, Moritz Saxenhofer,
Stephan Drewes, Mathias Schlegel,
Konrad M. Wanka, Raphael Frank, Sven
Klimpel, Felix von Blanckenhagen et al.**

Archives of Virology

Official Journal of the Virology
Division of the International Union of
Microbiological Societies

ISSN 0304-8608

Arch Virol

DOI 10.1007/s00705-016-2762-6



Your article is protected by copyright and all rights are held exclusively by Springer-Verlag Wien. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".

High genetic structuring of Tula hantavirus

Sabrina Schmidt¹ · Moritz Saxenhofer^{2,17} · Stephan Drewes¹ · Mathias Schlegel^{1,18} · Konrad M. Wanka¹ · Raphael Frank³ · Sven Klimpel³ · Felix von Blanckenhagen⁴ · Denny Maaz⁵ · Christiane Herden⁶ · Jona Freise⁷ · Ronny Wolf⁸ · Michael Stubbe⁹ · Peter Borkenhagen¹⁰ · Hermann Ansorge¹¹ · Jana A. Eccard¹² · Johannes Lang¹³ · Elsa Jourdain¹⁴ · Jens Jacob¹⁵ · Philippe Marianneau¹⁶ · Gerald Heckel^{2,17} · Rainer G. Ulrich¹ 

Received: 26 August 2015 / Accepted: 14 January 2016
© Springer-Verlag Wien 2016

Abstract Tula virus (TULV) is a vole-associated hantavirus with low or no pathogenicity to humans. In the present study, 686 common voles (*Microtus arvalis*), 249 field voles (*Microtus agrestis*) and 30 water voles (*Arvicola spec.*) were collected at 79 sites in Germany, Luxembourg and France and screened by RT-PCR and TULV-IgG ELISA. TULV-specific RNA and/or antibodies were detected at 43 of the sites, demonstrating a geographically widespread distribution of the virus in the studied area. The TULV prevalence in common voles (16.7 %) was higher than that in field voles (9.2 %) and water voles (10.0 %). Time series data at ten trapping sites showed evidence of a

lasting presence of TULV RNA within common vole populations for up to 34 months, although usually at low prevalence. Phylogenetic analysis demonstrated a strong genetic structuring of TULV sequences according to geography and independent of the rodent species, confirming the common vole as the preferential host, with spillover infections to co-occurring field and water voles. TULV phylogenetic clades showed a general association with evolutionary lineages in the common vole as assessed by mitochondrial DNA sequences on a large geographical scale, but with local-scale discrepancies in the contact areas.

Electronic supplementary material The online version of this article (doi:10.1007/s00705-016-2762-6) contains supplementary material, which is available to authorized users.

✉ Rainer G. Ulrich
rainer.ulrich@fli.bund.de

- ¹ Federal Research Institute for Animal Health, OIE Collaborating Centre for Zoonoses in Europe, Institute for Novel and Emerging Infectious Diseases, Friedrich-Loeffler-Institut, Suedufer 10, 17493 Greifswald, Insel Riems, Germany
- ² Computational and Molecular Population Genetics (CMPG), Institute of Ecology and Evolution, University of Bern, 3012 Bern, Switzerland
- ³ Goethe-University, Institute of Ecology, Evolution and Diversity, Senckenberg Biodiversity and Climate Research Centre, Senckenberg Gesellschaft für Naturforschung, 60438 Frankfurt am Main, Germany
- ⁴ RIFCON GmbH, 69493 Hirschberg, Germany
- ⁵ Institute for Parasitology and Tropical Veterinary Medicine, Freie Universität Berlin, 14163 Berlin, Germany
- ⁶ Institute for Veterinary Pathology, Justus-Liebig-Universität Gießen, 35392 Gießen, Germany

- ⁷ Task-Force Veterinärwesen, Fachbereich Schädlingsbekämpfung, Niedersächsisches Landesamt für Verbraucherschutz und Lebensmittelsicherheit, 26133 Oldenburg, Germany
- ⁸ Institute for Biology, University of Leipzig, 04103 Leipzig, Germany
- ⁹ Institute of Zoology, Martin-Luther-University Halle, 06099 Halle, Germany
- ¹⁰ Säugetierkundliche Arbeitsgemeinschaft Schleswig-Holstein, 24253 Probsteierhagen, Germany
- ¹¹ Senckenberg Museum of Natural History, 02826 Görlitz, Germany
- ¹² Institute for Biochemistry and Biology, Animal Ecology, University of Potsdam, 14469 Potsdam, Germany
- ¹³ Institut für Tierökologie und Naturbildung, Hauptstraße 30, 35321 Gonterskirchen, Germany
- ¹⁴ INRA, French National Institute for Agricultural Research, UR0346 Animal Epidemiology Unit, Saint-Genès Champanelle, France

Introduction

Hantaviruses (family *Bunyaviridae*, genus *Hantavirus*) were initially thought to be exclusively rodent-borne pathogens hosted by representatives of the families Muridae and Cricetidae [1]. The recent finding of novel hantaviruses in insectivores such as shrews and moles as well as bats raises questions about the origin and evolution of this group of viruses [2, 3].

Rodent-borne hantaviruses can cause two types of disease in humans, hantavirus cardiopulmonary syndrome (HCPS) and haemorrhagic fever with renal syndrome (HFRS). HCPS due to infection by New World hantaviruses, e.g., Sin Nombre virus (SNV) and Andes virus (ANDV), is associated with an average case fatality rate of about 40 % [4]. In Europe, hantaviruses are emerging pathogens with an increasing significance for human health [5], causing HFRS with differing case fatality rates, ranging from less than 1 % to 16 % [4, 6].

Hantaviruses associated with members of the vole genus *Microtus* have been detected in several parts of Europe, Asia and the North American continent. The North American California vole (*Microtus californicus*), meadow vole (*M. pennsylvanicus*) and prairie vole (*M. ochrogaster*) are known to harbour Isla Vista virus (ISLAV), Prospect Hill virus (PHV) and Bloodland Lake virus (BLLV), respectively [7]. In the Asian part of Russia and in China, three different hantaviruses have been detected in *Microtus* voles, with one species sometimes hosting more than one hantavirus: Khabarovsk virus (KHAV) was found in *M. fortis* and *M. maximowiczii*, Vladivostok virus (VLAV) in *M. fortis* and *M. oeconomus*, and Yuangjiang virus (YUJV) in *M. fortis* [8–12]. In Europe, two *Microtus*-associated hantaviruses have been described. Tula virus (TULV) was initially detected in common voles (*M. arvalis*) and sibling voles (*M. levis*, formerly *rossiaemeridionalis*) [13, 14]. A highly divergent TULV strain, designated as Adler virus, was recently discovered in common voles in Russia [15]. A second hantavirus, Tatenale virus, was found in field voles (*M. agrestis*) in England [16].

Like in all hantaviruses, the genome of TULV is partitioned into three segments: For TULV prototype strain

Moravia, the small (S) segment, 1,831 nucleotides (nt) in length, codes for the nucleocapsid (N) protein, the medium-sized (M) segment of 3,694 nt codes for two glycoproteins (Gn und Gc), and the large (L) segment of 6,541 nt codes for the viral RNA-dependent RNA polymerase [17–19]. In addition, the S segment of all vole-derived TULV strains contains an overlapping open reading frame (ORF) coding for a putative nonstructural protein (NSs) that has been shown to enhance survival of the virus during passaging in interferon-competent cells [20, 21].

There is little knowledge about the pathogenicity of TULV for humans. TULV-specific antibodies have been detected in healthy blood donors in the Czech Republic [19] and in German forestry workers, a potential risk group for hantavirus infections [22]. Furthermore, one HFRS patient from Germany had TULV-specific neutralizing antibodies [23]. In addition, TULV RNA was detected in EDTA blood of an acutely infected, immunocompromised patient in the Czech Republic [24].

Corresponding to the wide distribution range of its main host, the common vole, TULV-specific nucleic acid has been detected in several European countries, but these studies usually included only one or a few trapping sites in a specific region (see Ref. [14] and [25–30] and references therein). Whereas most hantaviruses are host-specific and natural spillover infections are only rarely reported, TULV has been molecularly detected in a wide variety of other Arvicolinae species: *M. levis*, *M. gregalis*, *M. subterraneus*, *M. agrestis*, *Lagurus lagurus*, and *Arvicola* spec. [13, 31–35]. Aside from common voles, the most TULV infections have been reported in field voles (*M. agrestis*) in some places even without the presence of the main host species or with a larger number of infected field voles compared to sympatric common voles. It was therefore speculated that rather than being spillover infected, field voles might represent another host species that can enable successful TULV replication [29, 34].

The evolution of hantaviruses in relation to their rodent hosts is controversial. Often, the divergence of hantaviruses and their rodent hosts is interpreted as a consequence of co-evolutionary processes. Alternatively, host switching and subsequent adaptation processes are thought to be a reason for the observed divergence patterns [2, 7, 34, 36–38]. Previous investigations were mainly focused only at the species level of the reservoir, but co-evolutionary processes may also occur at the level of evolutionary divergence within species [7, 29, 34, 35, 37]. In Central and Eastern Europe, four main evolutionary lineages of common voles have been identified with both autosomal and mitochondrial DNA markers: the Western, Central, Italian and Eastern lineages [39–41]. So far, nothing is known about potential co-divergence between TULV and evolutionary lineages of the common vole.

Here, we describe large-scale serological and RT-PCR TULV screening of three potential reservoir species from

¹⁵ Federal Research Centre for Cultivated Plants, Institute for Plant Protection in Horticulture and Forestry, Vertebrate Research, Julius Kühn-Institute, 48161 Münster, Germany

¹⁶ Virology Unit, Laboratory of Lyon, French Agency for Food, Environmental and Occupational Health and Safety (ANSES), 69364 Lyon, France

¹⁷ Swiss Institute of Bioinformatics, Genopode, 1015 Lausanne, Switzerland

¹⁸ Present Address: Seramun Diagnostica GmbH, 15754 Heidesee, Germany

Central and Western Europe, as well as phylogenetic analysis of TULV sequences and mitochondrial DNA sequences of common voles.

Materials and methods

Rodent trapping and dissection

Rodents were collected between 2004 and 2013 at 71 sites in 13 federal states of Germany, two sites in Luxembourg, and six sites in two administrative districts in France (Alsace and Midi-Pyrénées) in woodland and open field habitats (Fig. 1). Trapping of some of the animals has been described previously [42–45]. Rodent dissection was done according to standard protocols and resulted in the collection of heart, lung, liver, spleen, kidney, and brain, as well as tissue samples from the ear pinna and tail. Chest cavity fluid (CCF) was obtained by adding 1 ml of PBS [46]. For TULV-positive animals, TULV-negative animals

from selected sites, and individuals with inconclusive morphological species identification, mitochondrial *cytochrome b* (*cyt b*) sequences were determined [47]. For samples that did not allow a morphological identification, a sex-determination PCR was performed following established protocols [48, 49].

Nucleic acid isolation

RNA extraction was performed using a modified QIAzol protocol. Briefly, RNA extraction was performed using 1 ml of QIAzol® Lysis Reagent (QIAGEN, Hilden, Germany) and sterilized steel beads of 0.5 cm diameter (Isometall, Pleidelsheim, Germany). After tissue homogenization, 200 µl of chloroform was added, and the sample was mixed and thereafter centrifuged for 15 min at 4 °C and 11,900 g. The resulting supernatant was mixed with 500 µl of cold isopropanol (−20 °C), incubated at −20 °C for 20 min, and centrifuged again at 4 °C for 10 min at 11,900 g. The resulting pellet was washed once with 1 ml




Fig. 1 Location of the main trapping sites in Germany, Luxembourg and France. An additional trapping site (#78) was located outside the range of this map in the Aveyron region in southern France

of 75 % ethanol and thereafter with 1 ml 99.8 % ethanol, dried at 56 °C for no more than 5 min, and resuspended in DEPC-treated water.

DNA for *cyt b* analysis was obtained from tissue samples using conventional chloroform DNA extraction or tail lysis overnight [47, 50]. Briefly, for extraction, all tissue samples were incubated overnight at 56 °C and 400 rpm in 300 µl of lysis buffer containing 50 mM KCl, 10 mM Tris-HCl, pH 9.0, 0.45 % NP40, 0.45 % Tween 20 and 1 mg of proteinase K per ml.

RT-PCR and serology

Hantavirus RT-PCR assays of lung tissue samples were done according to previously described protocols for the PUUV/TULV S segment [51]. TULV-N-specific antibodies were detected in the CCF by IgG-ELISA using yeast-expressed, purified recombinant N protein of TULV strain Moravia [22, 34].

Statistical analysis

The significance of prevalence differences between sexes was investigated using the χ^2 test. Because of the low number of RNA-positive water voles, we used Fisher's exact test to evaluate the significance of prevalence differences between species.

Sequence determination and phylogenetic analysis

DNA sequencing was performed by the dideoxy-chain termination method using a BigDye Terminator v1.1 Kit (Applied Biosystems, Darmstadt, Germany) and Genetic Analyser 3130 and 3130xl sequencers (Applied Biosystems). When the direct sequencing approach failed, sequences were obtained after insertion of the RT-PCR product into the pCR®-TOPO®-vector and transformation of TOP10 cells according to the manufacturer's instructions (TOPO-TA-Cloning Kit, Invitrogen, Darmstadt, Germany). At least two plasmids per RT-PCR product were sequenced.

All generated data were subjected to a BLAST-mediated comparison of the novel sequences with sequences available in GenBank (<http://www.ncbi.nlm.nih.gov>). All TULV sequences were included in subsequent phylogenetic analysis, and identical sequences were excluded. For common vole lineage analysis, one to three representatives from each trapping site were chosen, with the exception of site #27, where mtDNA of 29 individuals was investigated. All TULV and mtDNA sequences were deposited in the GenBank database with accession numbers KU139527-KU139605 and KU139696-KU139816, respectively (Supplementary Tables 4 and 5).

Nucleotide sequences were aligned using the ClustalW method implemented in BioEdit v7.2.5 [52] and revised


Fig. 2 Maximum-likelihood (ML) phylogenetic tree based on TULV partial S-segment sequences (A) and association of TULV sequence clades with common vole evolutionary lineages (B). Novel sequences are labeled with sampling location, individual code, and host species. Published sequences obtained from GenBank are labeled with the accession number, location and host species. Bootstrap support for ML and posterior probabilities of Bayesian analyses are indicated for major branches only. — indicates bootstrap values <50; * indicates a different topology in Bayesian analysis. Clades Ia, Ib, II and III are major geographically coherent TULV clusters with adjacent distribution in Central Europe (see text). For GenBank accession numbers of the novel TULV sequences, see Supplementary Table 4

manually. In addition to the novel sequences obtained in this study, representative sequences were obtained from GenBank, and these are labeled with accession numbers in Fig. 2 and Fig. 3. The final datasets used for analysis contained 158 S-segment sequences of 255-bp length for the TULV S-segment and 148 sequences of 763 bp from the *cyt b* gene of *M. arvalis*. Reference sequences for *cyt b* analysis were chosen as described in Ref. [34] and [53]. The outgroup sequences included Puumala virus (PUUV), Prospect Hill virus (PHV), and Tatenale virus for TULV and *M. gregalis* for the *cyt b* dataset. The best nucleotide substitution model to fit each data-set was determined with jModeltest v2.1.6 [54]. The Tamura and Nei model with a gamma-distribution model of among-site rate heterogeneity and a proportion of invariable sites (TrN+G+I) had highest scores for the TULV and *cyt b* data according to the Bayesian Information Criterion (BIC). MEGA 6 [55] was used for phylogenetic tree reconstruction based on maximum-likelihood (ML) algorithms with 1000 bootstrap replications. Phylogenetic relationships were also inferred using the Bayesian method implemented in MrBayes v3.2.2 [56] on the CIPRES platform [57], employing individual nucleotide substitution rate priors for each dataset. Four independent analyses were done for TULV and *cyt b* data, comprising each 10^7 generations of Markov chain Monte Carlo chains, sampled every 10^3 generations with a burn-in fraction of 25 %. For both datasets, the average standard deviation of split frequencies was lower than 0.01 in every run, and the potential scale reduction factor was in the range of 0.99–1.01 for all parameters, indicating that parameter convergence had occurred. Consensus trees were drawn with FigTree v1.4.2 [58].

Results

Detection of TULV in common voles

To study the geographical distribution of TULV in its reservoir host, common voles were trapped in Germany, France and Luxembourg (Fig. 1, Table 1, Supplementary







Fig. 2 continued

Table 1). The 686 common voles that were collected originated from 51 sites in 47 districts of 13 federal states in Germany ($n = 654$), six sites in two administrative regions of France ($n = 26$) and one site in Luxembourg ($n = 6$).

A total of 115 (16.7 %) common voles had signs of a previous or ongoing TULV infection (Table 1). Hantavirus-specific RNA and IgG antibodies were detected in 107 of 685 (15.6 %) and 49 of 670 (7.3 %) animals, respectively. Nine common voles (0.9 %) were positive in ELISA only without any detectable RNA. When focusing on the 15 locations where ten or more animals had been captured, the molecular

prevalence ranged from 0 % to 37.5 % (mean: 14.2 %) and the seroprevalence ranged from 0 % to 19.0 % (mean: 5.3 %). TULV RNA could be detected in common voles from all three countries (Fig. 1, Supplementary Table 1). In contrast, TULV-specific antibodies were detected only in voles from Germany and Luxembourg. TULV-reactive antibodies were not found in any of the 26 animals from France, although TULV-RNA was detected in five animals originating from three trapping sites (Fig. 1, sites #75, #78, #79). For the RT-PCR-positive vole from site #79, no TULV sequence data were obtained after repeated attempts. Two out of six common voles from Luxembourg were RNA



labels specify the lineage and GenBank accession number. *M. gregalis* was used as an outgroup. Bootstrap support for ML and posterior probabilities of Bayesian analysis are indicated for major branches only. — indicates bootstrap values <50; *indicates a different topology in Bayesian analysis. For GenBank accession numbers of the *cytochrome b* sequences, see Supplementary Table 5

Consecutive trappings of voles at one site in France (#75) and nine of 14 sites in Germany (#09, #10, #27, #39, #40, #41, #44, #57, #65) revealed a continuous presence or re-appearance of TULV infections (Supplementary Table 2). The presence of TULV RNA was monitored and detected in the common vole populations for one month (sites #10, #27, #44, #65), several months

Table 1 Results of serological and RT-PCR investigations for *Microtus arvalis* (Marv), *Microtus agrestis* (Magr) and *Arvicola* spec. (Arv) collected in Germany, Luxembourg and France

Species	No. of trapping sites	Total no. of voles analyzed	Sex	Weight range (g)		No. positive by serology only / total no. analyzed	No. positive by PCR only / total no. analyzed	No. positive in both / total no. analyzed	Total no. positive by serology/ total no. analyzed	Total no. positive by RT-PCR / total no. analyzed	Total no. positive by serology and/ or RT-PCR	
				Male	Female							
<i>Marv</i>	58	686	320	367	5	48	9/670	67/685	40/668	49/670	107/685	115/686
							1.3 %	9.8 %	6.0 %	7.3 %	15.6 %	16.8 %
<i>Magr</i>	37	249	134	115	16	47	10/246	4/249	9/246	19/249	13/249	23/249
							4.1 %	1.6 %	3.7 %	7.6 %	5.2 %	9.2 %
<i>Arv</i>	6	30	14	16	21	132	2/29	1/30	0/30	2/29	1/30	3/30
							6.9 %	3.3 %	0 %	6.9 %	3.3 %	10.0 %

¹ no sequence for one animal

(sites #9, #41, #57 and #75), and up to several years (sites #39, #40).

Detection of TULV in field voles

Field voles were successfully collected at 37 locations in Germany (n = 249). In total, 23 of 249 (9.2 %) field voles were found to have TULV-specific RNA and/or antibodies (Table 1). Thirteen field voles from four trapping sites in three German federal states were RT-PCR positive (5.2 %), and virus-specific antibodies were also present in nine of them. Ten additional animals from five federal states were positive by ELISA, but no viral RNA was detectable (Table 1, Supplementary Table 1). When focusing on the seven trapping sites with 10 or more field voles captured, the molecular prevalence ranged from 0 % to 14.0 % (mean 2.3 %), and the seroprevalence ranged from 0 % to 16.0 % (mean: 3 %). Similar to common voles, TULV infections were detected more often in males (11.2 %) than in females (7.0 %) (Table 3), but this difference was not significant ($\chi^2 = 1.326$, df = 1, $p = 0.25$).

At five of six trapping sites (#1, #18, #39, #44, #65) with TULV-positive field voles, TULV RNA was also detected in sympatrically occurring common voles. At the remaining site (#42) all four common voles were TULV negative. Inversely, at five of 11 sites with TULV-infected common voles (sites # 1, #18, #39, #44, #65), sympatric field voles showed signs of TULV infection. However, at five of the six sites without TULV-positive field voles, only 1-3 animals were found to be infected. At the remaining site (#41), all 21 field voles that were trapped were TULV negative.

TULV infection in water voles

Water voles were successfully trapped at four sites in Germany (n = 26) and both sites in Luxembourg (n = 4; Table 1). Two of the 29 water voles were positive in TULV-ELISA, but no RNA could be amplified from those animals (Table 1, Supplementary Table 1). A TULV sequence could be obtained from one additional animal, although there was no reaction in the ELISA. All three TULV-positive water voles originated from Bavaria, Germany, but the two seropositive animals and the RT-PCR-positive animal were captured at different trapping sites about 12 km apart (Fig. 1, sites #44 and #51).

At five of the six trapping sites, water voles occurred sympatrically with common voles (sites #26, #73), field voles (site #51) or both species (sites #44 and #65). At site #44, detection of TULV RNA in the single trapped water vole was accompanied by detection of TULV-specific RNA and antibodies in common voles and TULV-specific antibodies in field voles (Supplementary Table 1). At site

Table 2 Sex differences in the detection of TULV-specific RNA and antibodies in the vole species *Microtus arvalis* (Marv), *Microtus agrestis* (Magr) and *Arvicola spec.* (Arv)

	No. positive / total no. analyzed (percentage)					
	Male			Female		
	Serology	RT-PCR	Serology and/or RT-PCR	Serology	RT-PCR	Serology and/or RT-PCR
Marv	28/310 (9.0)	57/319 (17.9)	61/320 (19.1)	21/361 (5.8)	50/366 (13.7)	54/367 (14.7)
Magr	13/132 (9.8)	8/134 (6.0)	15/134 (11.2)	6/114 (5.3)	5/115 (4.3)	8/115 (7.0)
Arv	1/13 (7.7)	0/14	1/14 (7.1)	1/16 (6.3)	1/16 (6.3)	2/16 (12.5)

Table 3 Mean nucleotide (A) and amino acid (B) sequence identity of the S segment and nucleocapsid protein sequences, respectively, between the TULV clades in Central and Western Europe (for definition of the clades, see Fig. 2; for their geographical origin see Fig. 4)

	Ia	Ib	II	III	IV
A					
Ia	94.2 %	-	-	-	-
Ib	87.9 %	91.5 %	-	-	-
II	82.8 %	81.9 %	91.6 %	-	-
III	81.8 %	79.6 %	79.7 %	88.1 %	-
IV	80.2 %	80.7 %	82.2 %	83.3 %	92.0 %
B					
Ia	99.8 %	98.4 %	94.5 %	89.9 %	89.9 %
Ib	-	97.7 %	94.3 %	89.2 %	89.1 %
II	-	-	99.0 %	91.1 %	91.9 %
III	-	-	-	97.5 %	94.7 %
IV	-	-	-	-	98.2 %

#51, TULV-specific antibodies were detected in two of the 17 water voles, but not in the three field voles trapped there. At two sites (#65, #73), TULV infections were detected in common voles or common and field voles, but not in the few water voles collected sympatrically. At the remaining two sites (#26, #72) none of the water voles and common voles showed signs of a TULV infection.

Sequence comparison and phylogenetic analysis of TULV sequences

To analyze the phylogenetic relationships between TULV strains, S segment sequences of all RT-PCR-positive voles were compared to existing TULV sequences from Germany and other European countries. The phylogenetic tree revealed a strong geographic structuring of TULV sequences that was independent of the rodent species of origin (Fig. 2). Main cluster I represents the majority of TULV sequences originating from Germany and comprises two parapatric sister clades: Ia (central and eastern

Germany) and Ib (northern, central and western Germany).

Main cluster II consists of novel sequences from the eastern part of Germany, published sequences from one site in this part of Germany [23], and prototype strain Lodz from Poland [59]. Cluster III contains sequences from southern Germany, Luxembourg, the Alsace region of France and a water-vole-derived TULV sequence from Switzerland [35]. TULV sequences from a trapping site in eastern Germany close to the Czech border (site #27), from Austria, and from the Czech Republic form cluster IV. The phylogenetic position of the novel sequence from southern France (site #78) could not be determined with confidence.

The intra-cluster proportion of variable sites ranged from 5.8 % to 11.9 % at the nt level and 0.2 % to 2.5 % at the amino acid level (Table 3). In contrast, the inter-cluster variability reached more than 20 % at the nt level and more than 10 % at the amino acid level, which exceeds the threshold level of 7 % established by the International Committee on Taxonomy of Viruses for hantavirus species definition based on the entire N protein sequence [60]. Sequence similarity at the nt and amino acid level was highest between the two subclades from Germany (Ia and Ib) and clade II from Eastern Germany and Poland (Tables 3A and 3B). The greatest differences were found between clusters Ia/Ib and III/IV, which differed on average by 8.9 % at the amino acid level and 20.3 % at the nt level (Tables 3A and 3B).

Co-segregation of TULV with evolutionary divergence in *M. arvalis*

To examine potential associations of TULV divergence with evolutionary divergence in the common vole, the *cyt b* sequences of selected animals were used for phylogenetic analysis together with reference sequences representing the evolutionary lineages Eastern, Central, Western and Italian (Fig. 3). The vast majority of *cyt b* sequences of common voles from Germany were identified as belonging to the Central lineage, and all investigated voles from France and




Fig. 4 Map showing the trapping sites of common voles infected with TULV of clusters Ia, Ib, II, III and IV and belonging to the evolutionary lineages Central, Eastern and Western

Luxembourg belonged to the Western lineage (Fig. 3 and Fig. 4). Interestingly, Central and Eastern evolutionary lineage *cyt b* sequences were identified in close proximity in the easternmost part of Germany, with sympatric occurrence of both lineages at site #27 (Fig. 4).

TULV clades showed a general association with certain evolutionary lineages of common voles. All animals harboring TULV of clusters Ia or Ib belonged to the Central lineage, except for one animal from Biebersdorf (site #18; Eastern lineage). TULV cluster III originated from common voles mainly of the Central lineage, with two from Luxembourg and one from France belonging to the Western lineage (sites #73 and #75). Sequences of cluster II are associated with the Eastern lineage but were also found in Central individuals trapped close to sites with Eastern lineage voles in western Brandenburg and southwestern Saxony (sites #28 and #79). TULV Cluster IV originated from Eastern-lineage hosts with one sequence from an animal of the Central lineage trapped at Cunnersdorf (site #27), Saxony. The TULV sequence from southern France originated from an animal of the Western lineage.

Discussion

Host specificity and spillover of TULV

In this study, we investigated three largely co-distributed vole species for the presence of TULV infections. The overall TULV prevalence for common voles (16.7 %) was higher than that for field voles (9.2 %) and water voles (10.0 %). In addition, the molecular prevalence for TULV differed significantly between species ($p < 0.001$; two-sided Fisher's exact test) and was again highest in common voles (15.6 %) compared to field voles (5.2 %) and water voles (3.3 %). Similarly, a previous study has shown a very low prevalence of TULV in water voles [35]. This may indicate a host preference of TULV for the common vole and the detection of TULV RNA in other vole species such as field and water voles being the result of spillover infections during the acute phase with active virus replication. In line with this assumption, almost all TULV-RNA-positive field voles were co-trapped with common voles or trapped at locations where TULV-RNA-positive

common voles were detected earlier and TULV sequences were relatively similar to those from common voles.

This observation of different prevalence in common voles versus field and water voles might also have been due to different interspecies interactions. Similar ecological characteristics and use of similar habitats can promote inter-specific contact between common voles and other species directly (fighting) or indirectly (feces and urine) [61, 62]. Common voles and field voles might coexist in the same habitat, while encounters between water voles and other microtine species usually lead to a consistent dominance of the water vole and exclusion of permanent *Microtus* populations during the breeding season [63]. The water voles trapped in our study seemed to be sympatric with *Microtus* species at five out of six trapping sites, but these two species (water voles and *Microtus* spp.) had not shared the same habitat for an extended period of time (Supplementary Table 2). This might explain why we could detect TULV RNA in water voles at only one location.

A host function of field voles was discussed in previous studies, as TULV RNA was detected at higher prevalence in field voles than in common voles or in multiple field vole individuals in the absence of common voles [29, 34]. In this study, the molecular prevalence in field voles was in general much lower than in common voles. Furthermore, 53 % of field voles with TULV-reactive IgG antibodies were TULV-RNA negative, which is indicative of an earlier infection with virus clearance, but TULV-RNA could be detected in 81.6 % of TULV-IgG-positive common voles, which is typical of persistent infection in the reservoir. Furthermore, almost all field voles from sites without sympatrically occurring common voles or with only TULV-negative common voles were TULV negative.

The overall results of this study support the notion that common voles act as the preferential host for TULV. Collectively, these data do not support the idea that field voles serve as reservoir hosts with an equal ability to support TULV replication and persistence. Additional studies are needed to determine whether field voles may temporally function as a reservoir host with low viral load.

TULV in common vole populations

TULV was detected at 39 of 62 (62.9 %) sites with common voles, but with rather low prevalence compared to other vole-associated hantaviruses. While our results settle well within the reported range of TULV RNA prevalence of 8 %-37 % [18, 64], other vole-associated hantaviruses such as PUUV have been reported to reach a molecular prevalence of 60 %-100 % in reservoir host populations [65, 66]. The TULV seroprevalence of no more than 19 % observed here is much lower than antibody prevalence of

up to 100 % reported for other Cricetidae- and Murinae-associated hantaviruses [66–71].

Our data suggest that – once in the common vole population – TULV seems to persist for months if not years. Persistence of hantaviruses in vole populations over several years has been shown for PUUV in bank voles. Distinct virus types were detectable over several years, independent of population density and virus prevalence [66]. Social behaviour, such as forming of colonies and kin clustering in winter, might contribute substantially to transmission and virus persistence, especially between closely related animals [72]. Our investigations have shown initial evidence of long-lasting presence of TULV in vole populations. Future systematic rodent-monitoring studies are needed to study potential intrinsic and extrinsic effects on TULV prevalence. Further studies should consider the potential oscillation of TULV prevalence in its reservoir according to season and long-term population dynamics [see ref. 66].

Discrepancy between serological and RT-PCR detection of TULV in the reservoir

This study shows a significant difference between seropositivity (49/670, 7.3 %) and molecular detection of TULV in common voles (105/685, 15.3 %) ($\chi^2 = 20.812$; $df = 1$; $p < 0.001$). One reason for this discrepancy might be a large number of acute TULV infections among common voles where no antibody response has been induced so far. This might be true in times of a high frequency of TULV transmission, perhaps during the peaks of the vole population. Alternatively, TULV might induce only a weak antibody response in the host, possibly due to an early innate immune response, as has been reported for *in vitro* investigations on the TULV-related PHV [73]. Furthermore, using a recombinant N protein originating from TULV strain Moravia for detection of other TULV strains in the ELISA might have caused a lower sensitivity of the IgG ELISA. Our sequence analysis (aa 225–307) including the hypervariable region of the N proteins [18] of members of TULV clades I to IV revealed sequence differences of up to 10.6 % at the amino acid level to the N protein of prototype strain Moravia (Supplementary Table 3). Additionally, the sensitivity of the ELISA might be influenced by the selection of the secondary antibody and setting of the cutoff value.

For nine adult common voles (weight, 18–36 g) TULV-reactive antibodies were found, but no viral nucleic acid, although TULV RNA was detected in other individuals from the same trapping sites and season. The detection of IgG antibodies in the absence of viral RNA may indicate virus clearance instead of a persistent infection. Similar

results were obtained in a study of PUUV in bank voles trapped in northern Sweden and for SNV in wild deer mice (*Peromyscus maniculatus*) [74, 75]. Furthermore, some studies of SEOV have suggested that, depending on the age of the reservoir upon infection, a persistent infection may not always be established [76]. Alternatively, this could be a sign of viral RNA load fluctuation during the course of infection.

In summary, our study showed a strong discrepancy between RT-PCR and serological detection of TULV infection in its reservoir. Future serological studies using antigens from different TULV strains should allow their role in the observed lower sensitivity of the ELISA compared to the RT-PCR to be tested. In addition, in-depth studies are needed to clarify the role of the rodent immune system in TULV infection and possible age effects.

Phylogeography of TULV and its reservoir

The detection of 79 novel TULV sequences from a large geographic area provided detailed insights into the high sequence variability and genetic structuring of TULV in Central Europe. Our study also allows an initial comparison of the phylogeography of the virus and its reservoir rodent host. In cluster II, the sequence of prototype strain Lodz from Poland is flanked by TULV sequences originating from common voles of the Eastern lineage, mostly found in countries located east of Germany, and four individuals of the Central lineage trapped at the eastern German border where Central and Eastern vole lineages meet [39, 40, 77–79]. In addition, our study shows the existence of distinct genetic clusters of TULV in close proximity to each other. This is consistent with the genetic isolation of common vole populations, even on a small geographic scale [40, 80, 81], and may indicate processes of virus-host adaptation.

In conclusion, this study shows that TULV is widespread in Central European common vole populations. TULV RNA was more frequently found in common voles than in field and water voles, confirming the common vole as the reservoir host and suggesting that infection of other vole species is most likely due to spillover. We readily detected TULV RNA in voles at different sites, but we were less successful in detecting specific antibodies. Although this could be a methodological problem, mechanisms leading to a reduced adaptive immune response cannot be excluded, and this provides an interesting target for further studies. The overall prevalence of TULV was not nearly as high as reported for the related PUUV in its bank vole reservoir. For further assessment of the potential involvement of population dynamics on TULV prevalence additional studies are needed. These monitoring studies would also have to consider potential consequences of

TULV infection on the fitness of the vole reservoir. The initial finding of large-scale associations of some TULV clades with different evolutionary lineages of common voles indicates the need for future studies in the contact areas to study potential (co-) evolutionary processes in more detail.

Acknowledgments The work in the laboratory of RGU was supported by the Robert Koch-Institut with funds from the German Ministry of Public Health (grant no. 1362/1-924 and 1362/1-980 to RGU), Deutsche Forschungsgemeinschaft (SPP 1596 “Ecology and Species Barriers in Emerging Viral Diseases”, UL 405/1-1 to RGU), and the EU grants APHAEA (grant no. 2811ERA117) and FP7-261504 EDENext and is catalogued by the EDENext Steering Committee as EDENext368 (<http://www.edenext.eu>). The contents of this publication are the sole responsibility of the authors and do not necessarily reflect the views of the European Commission. Collection of samples in Jeesser, Mecklenburg-Western Pomerania, were done in the frame of the UFOPLAN project 3709 41 401 granted to Jens Jacob. Moritz Saxenhofer was supported by grant 31003A-149585 from the Swiss National Science Foundation to Gerald Heckel. The technical assistance of Sybille Herzog, Jorge A. Encarnação, Séverine Murri, Daniel Nobach, Markus Eickmann, Manon Bourg, Kathrin Baumann, Samuel Bernstein, Thomas Büchner, Angele Breithaupt, Fabian Deutschens, Paul Dremsek, Ulrike Duve, Sylvia Ferguson, Henrike Gregersen, Bärbel Hammerschmidt, Dörte Kaufmann, Robert Klopffleisch, Nastasja G. Kratzmann, Christian Kretzschmar, Jens Lewitzki, Christina Maresch, Marc Mertens, Joachim Meyer, Ulrike M. Rosenfeld, Josephine Schlosser, Julia Schneider, Josephine Schröder, Kati Seve, Hanan Sheikh Ali, Jens P. Teifke, Franziska Thomas, Ute Wessels, Daniel Windolph, and Theres Wollny in rodent dissection and the collection and provision of rodents by Sabine Atger, Laurent Crespin, Anja Globig, Torsten Heidecke, Dietrich Heidecke, Anika Hellemann, Christian Imholt, Martin Kaatz, Joachim Kramer, Thilo Liesenjohann, Paul-Walter Löhr, Jennifer Maiano, Sébastien Massegli, Lutz Ohlmeyer, Angélique Pion, Daniela Reil, Dirk Steinhauer, Alexandre Teynier, Jörg Thiel, Cornelia Triebenbacher, Elise Vaumourin, Matthias Wenk and the RIFCON Field Study Team is kindly acknowledged. The authors would like to thank Martin H. Groschup and Noel Tordo for support, and Nicole Neumann for generation of Figures 1 and 4.

References

1. Plyusnin A, Elliott RM (2011) Bunyaviridae: molecular and cellular biology. Caister Academic Press, Norfolk
2. Henttonen H, Buchy P, Suputtamongkol Y, Jittapalpong S, Herbreteau V, Laakkonen J, Chaval Y, Galan M, Dobigny G, Charbonnel N, Michaux J, Cosson JF, Morand S, Hugot JP (2008) Recent discoveries of new hantaviruses widen their range and question their origins. *Ann N Y Acad Sci* 1149:84–89
3. Yanagihara R, Gu SH, Arai S, Kang HJ, Song JW (2014) Hantaviruses: rediscovery and new beginnings. *Virus Res* 187:6–14
4. Vaheri A, Strandin T, Hepojoki J, Sironen T, Henttonen H, Mäkelä S, Mustonen J (2013) Uncovering the mysteries of hantavirus infections. *Nat Rev Microbiol* 11:539–550
5. Heyman P, Ceianu CS, Christova I, Tordo N, Beersma M, João Alves M, Lundkvist A, Hukic M, Papa A, Tenorio A, Zelená H, Essbauer S, Visontai I, Golovljova I, Connell J, Nicoletti L, Esbroeck M Van, Gjeruldsen Dudman S, Aberle SW, Avšič-Županc T, Korukluoglu G, Nowakowska A, Klempa B, Ulrich

- RG, Bino S, Engler O, Opp M, Vaheri A (2011) A five-year perspective on the situation of haemorrhagic fever with renal syndrome and status of the hantavirus reservoirs in Europe, 2005–2010. *Euro Surveill* 16(36):pii=19961
6. Avsic-Zupanc T, Toney A, Anderson K, Chu YK, Schmaljohn C (1995) Genetic and antigenic properties of Dobrava virus: a unique member of the Hantavirus genus, family Bunyaviridae. *J Gen Virol* 76:2801–2808
7. Plyusnin A, Morzunov SP (2001) Virus evolution and genetic diversity of hantaviruses and their rodent hosts. *Curr Top Microbiol Immunol* 256:47–75
8. Cong M, Guo W, Wang J, Wang W, Zhou R, Li M, Zhang J, Zhang Y (2012) Genetic characteristics of hantaviruses carried by *Microtus maximowiczii* in Yakeshi of Inner Mongolia, China. *Zhonghua Liu Xing Bing Xue Za Zhi* 33:832–835
9. Hörling J, Chizhikov V, Lundkvist A, Jonsson M, Ivanov L, Dekonenko A, Niklasson B, Dzagurova T, Peters CJ, Tkachenko E, Nichol S (1996) Khabarovsk virus: a phylogenetically and serologically distinct hantavirus isolated from *Microtus fortis* trapped in far-east Russia. *J Gen Virol* 77:687–694
10. Plyusnina A, Laakkonen J, Niemimaa J, Nemirov K, Muruyeva G, Pohodiev B, Lundkvist Å, Vaheri A, Henttonen H, Vapalahti O, Plyusnin A (2008) Genetic analysis of hantaviruses carried by *Myodes* and *Microtus* rodents in Buryatia. *Virol J* 5:4
11. Zou Y, Wang JB, Gaowa HS, Yao LS, Hu G-W, Li MH, Chen HX, Plyusnin A, Shao R, Zhang YZ (2008) Isolation and genetic characterization of hantaviruses carried by *Microtus* voles in China. *J Med Virol* 80:680–688
12. Zou Y, Xiao QY, Dong X, Lv W, Zhang SP, Li MH, Plyusnin A, Zhang YZ (2008) Genetic analysis of hantaviruses carried by reed voles *Microtus fortis* in China. *Virus Res* 137:122–128
13. Plyusnin A, Vapalahti O, Lankinen H, Lehtväslaiho H, Apekina N, Myasnikov Y, Kallio-Kokko H, Henttonen H, Lundkvist A, Brummer-Korvenkontio M (1994) Tula virus: a newly detected hantavirus carried by European common voles. *J Virol* 68:7833–7839
14. Sibold C, Sparr S, Schulz A, Labuda M, Kozuch O, Lysý J, Krüger DH, Meisel H (1995) Genetic characterization of a new hantavirus detected in *Microtus arvalis* from Slovakia. *Virus Genes* 10:277–281
15. Tkachenko EA, Witkowski PT, Radosa L, Dzagurova TK, Okulova NM, Yunicheva YV, Vasilenko L, Morozov VG, Malkin GA, Krüger DH, Klempa B (2015) Adler hantavirus, a new genetic variant of Tula virus identified in Major's pine voles (*Microtus majori*) sampled in southern European Russia. *Infect Genet Evol* 29:156–163
16. Pounder KC, Begon M, Sironen T, Henttonen H, Watts PC, Voutilainen L, Vapalahti O, Klempa B, Fooks AR, McElhinney LM (2013) Novel hantavirus in field vole, United Kingdom. *Emerg Infect Dis* 19:673–675
17. Kukkonen SK, Vaheri A, Plyusnin A (1998) Completion of the Tula hantavirus genome sequence: properties of the L segment and heterogeneity found in the 3' termini of S and L genome RNAs. *J Gen Virol* 79:2615–2622
18. Plyusnin A, Cheng Y, Vapalahti O, Pejcoch M, Unar J, Jelinkova Z, Lehtväslaiho H, Lundkvist A, Vaheri A (1995) Genetic variation in Tula hantaviruses: sequence analysis of the S and M segments of strains from Central Europe. *Virus Res* 29:237–250
19. Vapalahti O, Lundkvist A, Kukkonen SK, Cheng Y, Gilljam M, Kanerva M, Manni T, Pejcoch M, Niemimaa J, Kaikusalo A, Henttonen H, Vaheri A, Plyusnin A (1996) Isolation and characterization of Tula virus, a distinct serotype in the genus Hantavirus, family Bunyaviridae. *J Gen Virol* 77:3063–3067
20. Jääskeläinen KM, Kaukinen P, Minskaya ES, Plyusnina A, Vapalahti O, Elliott RM, Weber F, Vaheri A, Plyusnin A (2007) Tula and Puumala hantavirus NSs ORFs are functional and the products inhibit activation of the interferon-beta promoter. *J Med Virol* 79:1527–1536
21. Jääskeläinen KM, Plyusnina A, Lundkvist A, Vaheri A, Plyusnin A (2008) Tula hantavirus isolate with the full-length ORF for nonstructural protein NSs survives for more consequent passages in interferon-competent cells than the isolate having truncated NSs ORF. *Virol J* 5:3
22. Mertens M, Hofmann J, Petraityte-Burneikiene R, Ziller M, Sasnauskas K, Friedrich R, Niederstrasser O, Krüger DH, Groschup MH, Petri E, Werdermann S, Ulrich RG (2011) Sero-prevalence study in forestry workers of a non-endemic region in eastern Germany reveals infections by Tula and Dobrava-Belgrade hantaviruses. *Med Microbiol Immunol* 200:263–268
23. Klempa B, Meisel H, Rath S, Bartel J, Ulrich R, Krüger DH (2003) Occurrence of renal and pulmonary syndrome in a region of northeast Germany where Tula hantavirus circulates. *J Clin Microbiol* 41:4894–4897
24. Zelená H, Mrázek J, Kuhn T (2013) Tula hantavirus infection in immunocompromised host, Czech Republic. *Emerg Infect Dis* 19:1873–1876
25. Klempa B, Radosa L, Krüger DH (2013) The broad spectrum of hantaviruses and their hosts in Central Europe. *Acta Virol* 57:130–137
26. Michalski A, Niemcewicz M, Bielawska-Drózd A, Nowakowska A, Gaweł J, Pitucha G, Joniec J, Zielonka K, Marciniak-Niemcewicz A, Kocik J (2014) Surveillance of hantaviruses in Poland: a study of animal reservoirs and human hantavirus disease in subcarpathia. *Vector Borne Zoonotic Dis* 14:514–522
27. Nikolic V, Stajkovic N, Stamenkovic G, Cekanac R, Marusic P, Siljic M, Gligic A, Stanojevic M (2014) Evidence of recombination in Tula virus strains from Serbia. *Infect Genet Evol* 21:472–478
28. Olsson GE, Leirs H, Henttonen H (2010) Hantaviruses and their hosts in Europe: reservoirs here and there, but not everywhere? *Vector-Borne Zoonotic Dis* 10:549–561
29. Scharninghausen JJ, Pfeffer M, Meyer H, Davis DS, Honeycutt RL, Faulde M (2002) Genetic evidence for Tula virus in *Microtus arvalis* and *Microtus agrestis* populations in Croatia. *Vector Borne Zoonotic Dis* 2:19–27
30. Schmidt S, Essbauer SS, Mayer-Scholl A, Poppert S, Schmidt-Chanasit J, Klempa B, Henning K, Schares G, Groschup MH, Spitzenberger F, Richter D, Heckel G, Ulrich RG (2014) Multiple infections of rodents with zoonotic pathogens in Austria. *Vector Borne Zoonotic Dis* 14:467–475
31. Iakimenko VV, Dekonenko AE, Mal'kova MG, Kuz'min IV, Tantsev AK, Dzagurova TK, Tkachenko EA (2000) The spread of hantaviruses in western Siberia. *Med Parazitol (Mosk)* 3:21–28
32. Korva M, Duh D, Puterle A, Trilar T, Županc TA (2009) First molecular evidence of Tula hantavirus in *Microtus voles* in Slovenia. *Virus Res* 144:318–322
33. Song JW, Gligic A, Yanagihara R (2002) Identification of Tula hantavirus in *Pitymys subterraneus* captured in the Cacak region of Serbia-Yugoslavia. *Int J Infect Dis* 6:31–36
34. Schmidt-Chanasit J, Essbauer S, Petraityte R, Yoshimatsu K, Tackmann K, Conraths FJ, Sasnauskas K, Arikawa J, Thomas A, Pfeffer M, Scharninghausen JJ, Spletstoeser W, Wenk M, Heckel G, Ulrich RG (2010) Extensive host sharing of central European Tula virus. *J Virol* 84:459–474
35. Schlegel M, Kindler E, Essbauer SS, Wolf R, Thiel J, Groschup MH, Heckel G, Oehme RM, Ulrich RG (2012) Tula virus infections in the Eurasian water vole in central Europe. *Vector Borne Zoonotic Dis* 12:503–513
36. Guo WP, Lin XD, Wang W, Tian JH, Cong ML, Zhang HL, Wang MR, Zhou RH, Wang JB, Li MH, Xu J, Holmes EC, Zhang YZ (2013) Phylogeny and origins of hantaviruses harbored by bats, insectivores, and rodents. *PLoS Pathogens* 9:e1003159

37. Nemirov K, Leirs H, Lundkvist A, Olsson GE (2010) Puumala hantavirus and *Myodes glareolus* in northern Europe: no evidence of co-divergence between genetic lineages of virus and host. *J Gen Virol* 91:1262–1274
38. Ramsden C, Melo FL, Figueiredo LM, Holmes EC, Zanutto PMA, VGDN Consortium (2008) High rates of molecular evolution in hantaviruses. *Mol Biol Evol* 25:1488–1492
39. Fink S, Excoffier L, Heckel G (2004) Mitochondrial gene diversity in the common vole *Microtus arvalis* shaped by historical divergence and local adaptations. *Mol Ecol* 13:3501–3514
40. Heckel G, Burri R, Fink S, Desmet JF, Excoffier L (2005) Genetic structure and colonization processes in European populations of the common vole, *Microtus arvalis*. *Evolution* 59:2231–2242
41. Lischer HEL, Excoffier L, Heckel G (2014) Ignoring heterozygous sites biases phylogenomic estimates of divergence times: implications for the evolutionary history of *Microtus* voles. *Mol Biol Evol* 31:817–831
42. Drexler JF, Corman VM, Müller MA, Maganga GD, Vallo P, Binger T, Gloza-Rausch F, Rasche A, Yordanov S, Seebens A, Oppong S, Sarkodie YA, Pongombo C, Lukashev AN, Schmidt-Chanasit J, Stöcker A, Carneiro AJB, Erbar S, Maisner A, Fronhoffs F, Buettner R, Kalko EKV, Kruppa T, Franke CR, Kallies R, Yandoko ERN, Herrler G, Reusken C, Hassanin A, Krüger DH, Matthee S, Ulrich RG, Leroy EM, Drosten C (2012) Bats host major mammalian paramyxoviruses. *Nat Commun* 3:796
43. Drexler JF, Corman VM, Müller MA, Lukashev AN, Gmyl A, Coutard B, Adam A, Ritz D, Leijten LM, van Riel D, Kallies R, Klose SM, Gloza-Rausch F, Binger T, Annan A, Adu-Sarkodie Y, Oppong S, Bourgarel M, Rupp D, Hoffmann B, Schlegel M, Kümmerer BM, Krüger DH, Schmidt-Chanasit J, Setién AA, Cottontail VM, Hemachudha T, Wacharapluesadee S, Osterrieder K, Bartenschlager R, Matthee S, Beer M, Kuiken T, Reusken C, Leroy EM, Ulrich RG, Drosten C (2013) Evidence for Novel Hepaciviruses in Rodents. *PLoS Pathogens* 9:e1003438
44. Guenther S, Grobbel M, Heidemanns K, Schlegel M, Ulrich RG, Ewers C, Wieler LH (2010) First insights into antimicrobial resistance among faecal *Escherichia coli* isolates from small wild mammals in rural areas. *Sci Total Environ* 408:3519–3522
45. Mayer-Scholl A, Hammerl J, Schmidt S, Ulrich R, Pfeffer M, Woll D, Scholz H, Thomas A, Nöckler K (2014) *Leptospira* spp. in Rodents and Shrews in Germany. *Int J Environ Res Public Health* 11:7562–7574
46. Ulrich RG, Schmidt-Chanasit J, Schlegel M, Jacob J, Pelz H-J, Mertens M, Wenk M, Büchner T, Masur D, Seve K, Groschup MH, Gerstengarbe F-W, Pfeffer M, Oehme R, Wegener W, Bemann M, Ohlmeyer L, Wolf R, Zoller H, Koch J, Brockmann S, Heckel G, Essbauer SS (2008) Network “Rodent-borne pathogens” in Germany: longitudinal studies on the geographical distribution and prevalence of hantavirus infections. *Parasitol Res* 103:121–129
47. Schlegel M, Ali HS, Stieger N, Groschup MH, Wolf R, Ulrich RG (2012) Molecular identification of small mammal species using novel cytochrome b gene-derived degenerated primers. *Biochem Genet* 50:440–447
48. Aasen E, Medrano JF (1990) Amplification of the ZFY and ZFX genes for sex identification in humans, cattle, sheep and goats. *Biotechnology (N Y)* 8:1279–1281
49. Bryja J, Konecny A (2003) Fast sex identification in wild mammals using PCR amplification of the Sry gene. *Folia Zoologica* 52:269–274
50. Schlegel M, Sheikh Ali H, Keller M, Groschup M, Ulrich R (2013) Molekulare Artbestimmung von Kleinsäugetern. *LabLoeffler* 7:16–17
51. Essbauer SS, Schmidt J, Conraths FJ, Friedrich R, Koch J, Hautmann W, Pfeffer M, Wölfel R, Finke J, Dobler G, Ulrich R (2006) A new Puumala hantavirus subtype in rodents associated with an outbreak of Nephropathia epidemica in South-East Germany in 2004. *Epidemiol Infect* 134:1333–1344
52. Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acid Symp Ser* 41:95–98
53. Braaker S, Heckel G (2009) Transalpine colonisation and partial phylogeographic erosion by dispersal in the common vole *Microtus arvalis*. *Mol Ecol* 18:2518–2531
54. Darriba D, Taboada GL, Doallo R, Posada D (2012) jModelTest 2: more models, new heuristics and parallel computing. *Nat Methods* 9:772
55. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 30:2725–2729
56. Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biol* 61:539–542
57. Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: Gateway Computing Environments Workshop (GCE), 2010, pp 1–8
58. Rambaut A (2012) FigTree v1. 4. Molecular evolution, phylogenetics and epidemiology. University of Edinburgh, Institute of Evolutionary Biology, Edinburgh
59. Song JW, Baek LJ, Song KJ, Skrok A, Markowski J, Bratosiewicz-Wasik J, Kordek R, Liberski PP, Yanagihara R (2004) Characterization of Tula virus from common voles (*Microtus arvalis*) in Poland: evidence for geographic-specific phylogenetic clustering. *Virus Genes* 29:239–247
60. Plyusnin A, Beaty BJ, Elliott RM, Goldbach R, Kormelink R, Lundkvist A, Schmaljohn CS, Tesh RB (2011) Bunyaviridae. In: King AMQ, Adams MJ, Carstens EB, Lefkowitz EJ (eds) *Virus taxonomy: ninth report of the international committee on taxonomy of viruses*. Elsevier Inc., San Diego, pp 725–741
61. Jacob J, Manson P, Barfknecht R, Fredricks T (2014) Common vole (*Microtus arvalis*) ecology and management: implications for risk assessment of plant protection products. *Pest Manag Sci* 70:869–878
62. Yletyinen S, Norrdahl K (2008) Habitat use of field voles (*Microtus agrestis*) in wide and narrow buffer zones. *Agr Ecosyst Environ* 123:194–200
63. Myllymäki A (1977) Interactions between the field vole *Microtus agrestis* and its microtine competitors in Central-Scandinavian populations. *Oikos* 29:570–580
64. Jakab F, Horváth G, Ferenczi E, Sebők J, Szűcs G (2008) First detection of Tula hantaviruses in *Microtus arvalis* voles in Hungary. *Arch Virol* 153:2093–2096
65. Essbauer SS, Schmidt-Chanasit J, Madeja EL, Wegener W, Friedrich R, Petraityte R, Sasnauskas K, Jacob J, Koch J, Dobler G, Conraths FJ, Pfeffer M, Pitra C, Ulrich RG (2007) Nephropathia epidemica in metropolitan area, Germany. *Emerg Infect Dis* 13:1271–1273
66. Weber de Melo V, Sheikh Ali H, Freise J, Kühnert D, Essbauer S, Mertens M, Wanka KM, Drewes S, Ulrich RG, Heckel G (2015) Spatiotemporal dynamics of Puumala hantavirus associated with its rodent host, *Myodes glareolus*. *Evol Appl* 8:545–559
67. Engelthaler DM, Levy CE, Fink TM, Tanda D, Davis T (1998) Short report: Decrease in seroprevalence of antibodies to hantavirus in rodents from 1993–1994 hantavirus pulmonary syndrome case sites. *Am J Trop Med Hyg* 58:737–738

68. Garanina SB, Platonov AE, Zhuravlev VI, Murashkina AN, Yakimenko VV, Korneev AG, Shipulin GA (2009) Genetic diversity and geographic distribution of hantaviruses in Russia. *Zoonoses Public Health* 56:297–309
69. Klein TA, Kang HJ, Gu SH, Moon S, Shim SH, Park YM, Lee SY, Kim HC, Chong ST, O'Guinn M, Lee JS, Turell MJ, Song JW (2011) Hantaan virus surveillance targeting small mammals at Dagmar North Training Area, Gyeonggi Province, Republic of Korea, 2001–2005. *J Vector Ecol* 36:373–381
70. Tadin A, Turk N, Korva M, Margaletić J, Beck R, Vucelja M, Habuš J, Svoboda P, Županc TA, Henttonen H, Markotić A (2012) Multiple co-infections of rodents with hantaviruses, *Leptospira*, and *Babesia* in Croatia. *Vector Borne Zoonotic Dis* 12:388–392
71. Yanagihara R, Daum CA, Lee PW, Baek LJ, Amyx HL, Gajdusek DC, Gibbs CJ Jr (1987) Serological survey of Prospect Hill virus infection in indigenous wild rodents in the USA. *Trans R Soc Trop Med Hyg* 81:42–45
72. Deter J, Chaval Y, Galan M, Gauffre B, Morand S, Henttonen H, Laakkonen J, Voutilainen L, Charbonnel N, Cosson JF (2008) Kinship, dispersal and hantavirus transmission in bank and common voles. *Arch Virol* 153:435–444
73. Handke W, Oelschlegel R, Franke R, Krüger DH, Rang A (2009) Hantaan virus triggers TLR3-dependent innate immune responses. *J Immunol* 182:2849–2858
74. Alexeyev OA, Ahlm C, Elgh F, Aava B, Palo T, Settergren B, Tärnvik A, Wadell G, Juto P (1998) A minority of seropositive wild bank voles (*Clethrionomys glareolus*) show evidence of current Puumala virus infection. *Epidemiol Infect* 121:419–425
75. Kuenzi AJ, Douglass RJ, Bond CW, Calisher CH, Mills JN (2005) Long-term dynamics of Sin Nombre viral RNA and antibody in deer mice in Montana. *J Wildl Dis* 41:473–481
76. Meyer BJ, Schmaljohn CS (2000) Persistent hantavirus infections: characteristics and mechanisms. *Trends Microbiol* 8:61–67
77. Beysard M, Heckel G (2014) Structure and dynamics of hybrid zones at different stages of speciation in the common vole (*Microtus arvalis*). *Mol Ecol* 23:673–687
78. Fischer MC, Foll M, Heckel G, Excoffier L (2014) Continental-scale footprint of balancing and positive selection in a small rodent (*Microtus arvalis*). *PLoS One* 9:e112332
79. Martinkova M, Barnett R, Cucchi T, Struchen R, Pascal M, Pascal M, Fischer MC, Higham T, Brace S, Ho SYW, Quere JP, O'Higgins J, Excoffier L, Heckel G, Hoelzel AR, Dobney KM, Searle JB (2013) Divergent evolutionary processes associated with colonization of offshore islands. *Mol Ecol* 22:5205–5220
80. Beysard M, Krebs-Wheaton R, Heckel G (2015) Tracing reinforcement through asymmetrical partner preference in the European common vole *Microtus arvalis*. *BMC Evol Biol* 15:170
81. Schweizer M, Excoffier L, Heckel G (2007) Fine-scale genetic structure and dispersal patterns in the common vole *Microtus arvalis*. *Mol Ecol* 16:2463–2473